# CORRECTION Open Access



# Correction: Neurotensin promotes the progression of malignant glioma through NTSR1 and impacts the prognosis of glioma patients

Qing Ouyang<sup>1,2†</sup>, Xueyang Gong<sup>2†</sup>, Hualiang Xiao<sup>3†</sup>, Ji Zhou<sup>1</sup>, Minhui Xu<sup>1</sup>, Yun Dai<sup>4</sup>, Lunshan Xu<sup>1</sup>, Hua Feng<sup>5</sup>, Hongjuan Cui<sup>2\*</sup> and Liang Yi<sup>1\*</sup>

# Correction: Mol Cancer 14, 21 (2015) https://doi.org/10.1186/s12943-015-0290-8

Following publication of the original article [1], the authors subsequently identified an error in Figure 4A. The images representing U87 cells were taken from GL261 cells by mistake. The corrected Figure 4 is now shown in this correction. The authors confirm that the conclusions of this article are not affected, and sincerely apologize for this error and any inconvenience that may have caused. The correct figure is as follows.

## Author details

<sup>1</sup>Department of Neurosurgery, Daping Hospital, Third Military Medical University, Chongqing, China. <sup>2</sup>State Key Laboratory of Silkworm Genome Biology, Institute of Sericulture and Systems Biology, Southwest University, Chongqing, China. <sup>3</sup>Department of Pathology, Daping Hospital, Third Military Medical University, Chongqing, China. <sup>4</sup>Division of Hematology/Oncology, Department of Medicine, Virginia Commonwealth University, Richmond, VA,

The original article can be found online at https://doi.org/10.1186/s12943-015-0290-8.

<sup>†</sup>Qing Ouyang, Xueyang Gong and Hualiang Xiao contributed equally to this work.

\*Correspondence: hongjuan.cui@gmail.com; happyape@126.com

USA. <sup>5</sup>Department of Neurosurgery, Southwest Hospital, Third Military Medical University, Chongqing, China.

Published online: 10 August 2022

### Reference

Ouyang Q, Gong X, Xiao H, et al. Neurotensin promotes the progression of malignant glioma through NTSR1 and impacts the prognosis of glioma patients. Mol Cancer. 2015;14:21. https://doi.org/10.1186/s12943-015-0290-8.




© The Author(s) 2022. **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit http://creativeccommons.org/licenses/by/4.0/. The Creative Commons Public Domain Dedication waiver (http://creativeccommons.org/publicdomain/zero/1.0/) applies to the data made available in this article, unless otherwise stated in a credit line to the data.

 $<sup>\</sup>overline{\ }$  Department of Neurosurgery, Daping Hospital, Third Military Medical University, Chongqing, China

<sup>&</sup>lt;sup>2</sup> State Key Laboratory of Silkworm Genome Biology, Institute of Sericulture and Systems Biology, Southwest University, Chongqing, China Full list of author information is available at the end of the article

Ouyang et al. Molecular Cancer (2022) 21:162 Page 2 of 2



**Fig. 4** NTS/NTSR1 boosted the migration capacity and invasiveness of glioma cells. **A**, Transwell invasion assay showed that the ability of GL261 and U87 glioma cells to invade across the matrigel and membrane. **B**, The proportion of invasive GL261 and U87 glioma cells in all the experimental groups in the transwell experiments. \*p < 0.01 vs. the control group using a two-tailed t test, & p < 0.01 vs. the NTS group using a two-tailed t test. \*p < 0.01 vs. the NTS group using a two-tailed t test. \*p < 0.01 vs. the NTS group using a two-tailed t test. \*p < 0.01 vs. the percentage of wound closure by the cells was quantified in the different groups at 36 hours and 72 hours. &p < 0.01 vs. the NTS group using a two-tailed t test. \*p < 0.01 vs. the control group using a two-tailed t test, \*p < 0.01 vs. the NTS group using a two-tailed t test.