

CORRECTION

Open Access



Correction: Gamabufotalin, a bufadienolide compound from toad venom, suppresses COX-2 expression through targeting IKK β /NF- κ B signaling pathway in lung cancer cells

Zhenlong Yu^{1†}, Wei Guo^{1†}, Xiaochi Ma^{1*}, Baojing Zhang¹, Peipei Dong¹, Lin Huang¹, Xiuli Wang¹, Chao Wang¹, Xiaokui Huo¹, Wendan Yu¹, Canhui Yi¹, Yao Xiao¹, Wenjing Yang¹, Yu Qin¹, Yuhui Yuan¹, Songshu Meng¹, Quentin Liu^{1,2} and Wuguo Deng^{1,2*}

Correction: *Mol Cancer* 13, 203 (2014)

<https://doi.org/10.1186/1476-4598-13-203>

Unfortunately, the original version of this article [1] contained some errors. The words “binding” has been spelt incorrectly in the fifth title of result section and legend of Fig. 5. And the image of GAPDH (Fig. 6D) has been mistakenly uploaded. The mistakes did not affect any correctness of our results or discussion. The corrected sentence Fig. 6 was provided here. We regret any inconvenience that caused.

“CS-6 inhibited NF- κ B and p300 translocation and binding to COX-2 promoter”. “Figure 5 CS-6 inhibited NF- κ B and p300 translocation and their binding to COX-2 promoter”.

[†]Zhenlong Yu and Wei Guo these authors contributed equally to this work.

The online version of the original article can be found at <https://doi.org/10.1186/1476-4598-13-203>

*Correspondence:

Xiaochi Ma
maxc1978@163.com

Wuguo Deng
dengwg@hotmail.com

¹Institute of Cancer Stem Cell, College of Pharmacy, Dalian Medical University, Lvshun South Road No 9, 116044 Dalian, China

²State Key Laboratory of Oncology in South China, Collaborative Innovation Center of Cancer Medicine, Sun Yat-sen University Cancer Center, Guangzhou, China



© The Author(s) 2023. **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated in a credit line to the data.




Fig. 6 CS-6 inhibited the phosphorylation and activation of IKK β . (D) A549 cells were treated with CS-6 (50 nM) after pretreatment with CB (50 nM). The IKK β , and p-IKK β proteins were analyzed by Western blotting.

Published online: 02 September 2023

signaling pathway in lung cancer cells. *Mol Cancer* **13**, 203 (2014). <https://doi.org/10.1186/1476-4598-13-203>

References

1. Yu, Z., Guo, W., Ma, X. et al Gamabufotalin, a bufadienolide compound from toad venom, suppresses COX-2 expression through targeting IKK β /NF- κ B

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.