

RESEARCH

Open Access



Histone demethylase KDM4D promotes gastrointestinal stromal tumor progression through HIF1 β /VEGFA signalling

Fuqing Hu, Haijie Li, Lu Liu, Feng Xu, Senyan Lai, Xuelai Luo, Junbo Hu and Xi Yang* 

Abstract

Background: Gastrointestinal stromal tumour (GIST) is the most common soft tissue sarcoma. The identification of the molecular mechanisms regulating GIST progression is vital for its treatment and prevention. Increasing reports have demonstrated that epigenetic alterations play critical roles in GIST development. However, the role of the histone demethylase KDM4D in GIST progression is poorly understood.

Methods: In clinically matched GIST tissues, KDM4D protein levels were measured by Western blot and immunohistochemical (IHC) staining. KDM4D mRNA levels were examined by quantitative real-time PCR (qRT-PCR). Bioinformatics analysis was used to examine KDM4D expression. The biological effects of KDM4D were investigated in vitro using CCK-8, BrdU/PI, wound healing, colony formation, tube formation and Transwell assays and in vivo using a xenograft mice model. Luciferase assays were used to assess regulation of HIF1 β gene promoter activity by KDM4D. ChIP assays were performed to assess KDM4D, H3K36me3 and H3K9me3 occupancy on the HIF1 β gene promoter.

Results: We observed a significant upregulation of KDM4D in GIST tissue compared with matched normal tissue and further explored the oncogenic function of KDM4D both in vitro and in vivo. Furthermore, we demonstrated that KDM4D directly interacted with the HIF1 β gene promoter and regulated its activity, promoting tumour angiogenesis and GIST progression both in vitro and in vivo. Finally, we demonstrated that KDM4D transcriptionally activates HIF1 β expression via H3K9me3 and H3K36me3 demethylation at the promoter region.

Conclusions: Our findings reveal the important roles of the KDM4D/HIF1 β /VEGFA signalling pathway in GIST progression, and this pathway may act as a potential therapeutic target for GIST patients.

Keywords: GIST - KDM4D, Proliferation, Migration, Angiogenesis

Background

Gastrointestinal stromal tumour (GIST) is the most common soft tissue sarcoma and often localizes to the gastrointestinal tract [1, 2]. Currently, the majority of studies indicate that GISTs originate from the mesenchymal pacemaker cells of the gastrointestinal tract known as the interstitial cells of Cajal (ICCs) that harbour multi-oncogenic mutations, such as KIT and PDGFRA [3, 4]. Increasing evidence has demonstrated that those oncogenes play a critical role in GIST tumourigenesis, proliferation, and metastasis. Given the important role

of oncogenes in GIST progression, molecular targeted drugs (imatinib) have been employed to cure GISTs harbouring mutant KIT or PDGFRA [5]. Although targeted drugs have revolutionized the treatment of GIST, a significant number of GIST patients experience recurrence within two years due to resistance [6, 7]. In addition, there is no promising treatment for wild-type KIT/PDGFRA GISTs [8]. Thus, to develop novel therapeutic strategies, further understanding of the molecular mechanisms of GISTs is crucial.

Recently, numerous studies have implied that epigenetic alterations play critical roles in a wide range of tumours [9, 10]. Previous studies have also demonstrated that epigenetic alterations are responsible for GIST development

* Correspondence: yx86ali@hust.edu.cn

Cancer Research Institute, Tongji Hospital, Huazhong University of Science and Technology, Wuhan, China



[11]. Both DNA hypomethylation and DNA hypermethylation are reported to be closely related to GIST progression. Igarashi S. reported that LINE-1 methylation was associated with malignant GIST profiles and poor prognosis. In addition, more genes are methylated in advanced GIST compared with benign GIST [12]. More important, DNA methylation is associated with aggressive clinical characteristics, strongly indicating that DNA methylation is involved in GIST progression and may act as a novel treatment approach for GIST patients [13]. In addition to DNA methylation, histone methylation is another major epigenetic modification that is a reversible process. Previous studies have implied that changes in histone methylation could lead to gene activation or repression and effect tumour progression [14, 15]. In GIST, histone H2AX is a direct mediator of gastrointestinal stromal tumour cell apoptosis upon treatment with imatinib mesylate [16]. Histones can be modified by methylation and demethylation. Numerous demethylases are involved in diverse tumour development [17]. For example, KDM4 family members demethylate different sites of histones to activate or suppress gene expression [18–20]. However, the potential role of demethylases in GIST remains largely unknown. Importantly, the molecular mechanisms by which demethylases regulate GIST progression remain unclear.

Herein, we demonstrate that KDM4D mRNA and protein levels are upregulated in GIST compared with the matched normal tissues. Further, KDM4D overexpression strongly promotes GIST cell proliferation, migration, invasion and tumour angiogenesis. In contrast, silencing KDM4D reduced cell proliferation, migration, invasion and tumour angiogenesis. More importantly, these biologic effects mediated by KDM4D might be dependent on the Hif1 β /VEGFA signalling pathway. These findings highlight novel evidence for a functional link between demethylase and GIST development. We believe that the KDM4D/Hif1 β /VEGFA signalling pathway may act as a potential therapeutic target for GIST patients.

Methods

Cell lines and reagents

GIST882 cells, a kind gift of Jonathan Fletcher (Dana-Farber Cancer Institute, Boston, MA), were derived from a GIST patient with a homozygous missense mutation in KIT exon 13 (K642E). GIST-T1 cells obtained from Biowit Technologies (Shenzhen, China) were derived from a GIST patient with an in-frame deletion of 57 nucleotides in KIT exon 11 (V560Y579del). Both cell lines used in this study were cultured at 37 °C with 5% CO₂ in DMEM supplemented with 1% penicillin-streptomycin and 10% fetal bovine serum. Antibodies to KDM4D (ab93694) and VEGFA (ab1316) were purchased from Abcam (Cambridge, MA, USA). HIF1a (#36169), H3K36me3 (#4909) and H3K9me3 (#13969) antibodies were obtained from Cell Signaling

Technology (Danvers, MA, USA). Antibodies to GAPDH (sc-47,724), HIF1 β (sc-17,811), Histone3 (sc-517,576) and mouse IgG (sc-69,786) were purchased from Santa Cruz Biotechnology (Santa Cruz, CA, USA). The human VEGFA ELISA Kit (EK0539) was obtained from Boster Biological Technology (Wuhan, China).

Immunohistochemistry

Human GISTs samples were obtained from Tongji Hospital. For immunohistochemistry, formalin-fixed paraffin-embedded tissues were cut into 4-mm sections. Then, slides were chosen to stain with antibodies against KDM4D, VEGFA and CD31, separately. Two experienced pathologists examined immunostaining results. The intensity stained scores were evaluated according to the intensity of the nucleic or cytoplasmic staining (no staining = 0, weak staining = 1, moderate staining = 2, strong staining = 3). Finally, the IHC score was generated by combining the intensity scores with the percentage of positively stained cells.

Quantitative real-time PCR (qRT-PCR)

Total RNA was extracted using TRIzol Reagent (Invitrogen) and reverse transcribed using SuperScript II Reverse Transcriptase (Invitrogen) according to the manufacturer's instructions. Quantitative PCR was performed using the ABI 7300 real-time PCR system (Applied Biosystems) with KDM4D and GAPDH specific primers. The fold-change in KDM4D expression was calculated using the $2^{-\Delta\Delta CT}$ method, and GAPDH mRNA levels served as the control. The sense primer for KDM4D was 5'-GGGCAGGGGTGTTTACTCAAT-3', the antisense primer for KDM4D was 5'-IGTTTGCCAAATGGCGATACT-3'. The sense primer for GAPDH was 5'-ACCA CAGTCCATGCCATCAC-3', the antisense primer for GAPDH was 5'-TCCACCACCCTGTTGCTG TA-3'.

Western blot

Total cell protein was isolated using NP40 lysates with protease and phosphatase inhibitors. Protein samples were separated by SDS-PAGE and then transferred to PVDF membranes. Blots were incubated with primary antibodies overnight at 4 °C. Then, blots were incubated with the corresponding HRP-conjugated secondary antibodies. Immunoreactive bands were examined with ECL reagents (Thermo Scientific).

Cell viability assay

Cells were seeded into 96-well plates. Overnight, the medium was exchanged with 100 μ l of medium supplemented with 10 μ l CCK8. The plates were incubated for 2 h. Afterwards, absorbance at 450 nm was measured to obtain OD value. Similarly, other time points (24, 48, 72 and 96 h) were also assessed according to the procedures.

Soft agar colony formation assay

Briefly, 1% noble agar was added to the bottom of 6-well plates. After the layer of agar solidified, 500 cells were seeded into the 6-well plates. The time required for adequate colony formation varies for each cell line and is typically approximately two weeks. The medium was replaced every three days. After approximately two weeks, photographs of colonies were obtained using a microscope.

ChIP assay

Chromatin immunoprecipitation assays were performed as described previously [20]. The primers used for CHIP assays are provided below: HIF1 β upstream sense, 5'-CGCTCTTGTTGCCAGACTGG-3'; HIF1 β downstream sense, 5'-TCTGTAATCCCAGCACTTTGGG-3'.

Luciferase reporter assay

Cells were co-transfected with 0.2 μ g of HIF1 β promoter-luciferase reporter plasmid (PGL3-HIF1 β promoter) and 10 ng Renilla luciferase control vector. Firefly luciferase activity was detected using dual-luciferase Reporter Assays (Promega) 48 h after transfection and normalized to Renilla luciferase activity. PGL3-HIF1 β promoter was generated using the following primers. The sense primer for the HIF1 β promoter was 5'-CTTTGTGATCCGCCCTCCTTG-3', and the anti-sense primer for the HIF1 β promoter was 5'-AGTAGGCGGAGTCAACACAC-3'.

BrdU/PI assay

BrdU/PI assays were performed as described previously [21]. Briefly, cells were cultured in a 6-well plate overnight followed by a 20-min incubation with 10 μ g/ml BrdU. Then, cells were washed twice with PBS and fixed with 70% ethanol at -20 °C overnight.

Next, cells were denatured in 2 N HCL for 45 min and stained with FITC secondary antibody for 1 h at room temperature. Cells were further incubated with 40 g/ml RNase A and 200 g/ml PI for 30 min and finally analysed by flow cytometry.

Endothelial cell tube formation assay

GIST cells were treated as indicated overnight in DMEM supplemented without serum. Then, 20,000 HUVECs cells per well incubated with GIST cell-conditioned media were seeded into the Matrigel in 96-well plates. Microscopic images of tube formation were obtained, and the number of branches in the formed HUVEC tubes was assessed.

Enzyme-linked immunosorbent assay (ELISA)

GISTs cells were treated as indicated overnight in DMEM supplemented without serum. Overnight, cell-conditioned medium was collected for VEGFA-ELISA analysis according

to the manufacturer's instructions. The OD values were obtained based on absorbance at 470 nm.

Wound healing and Transwell assays

For the wound-healing assay, GISTs cells were seeded on six-well plates. When 95% confluence was achieved, the cell monolayer was gently scratched using a 200- μ m sterile plastic pipette tip. Then, the wound was photographed. After 24 h, the healing wound was photographed. For Transwell migration or invasion assays, 4×10^4 cells suspended in medium without serum were seeded in the upper chamber membranes coated without/with Matrigel (BD Biosciences). Then, 600 μ l medium with 10% FBS was added to the lower chamber. After 24 h, the underside of the membrane was fixed for 30 min and stained with 0.1% crystal violet. The inner side of the membrane was wiped with a cotton swab. Then, cells were quantified under a microscope.

Mouse xenograft tumour assay

All mice experiments were approved by the Animal Care and Use Committee of Tongji Hospital. Four-week Balb/c nude mice were obtained from Beijing Huafukang Bioscience, and 12 mice were randomly divided into 2 groups. We resuspended GIST882 cells stably expressing vector or KDM4D in PBS, mixed the cells with Matrigel in a 1:1 ratio and injected the suspension into the subcutaneous right groin tissue of Balb/c nude mice. After these mice developed a visible tumour mass, tumour volumes were calculated as (length \times width \times width)/2. Thirty days after subcutaneous injection, mice were sacrificed. Of note, for the mouse xenograft tumor model from ShNC cells and ShKDM4D cells, mice were sacrificed after subcutaneous injection 25d.


Statistical analysis

Statistical analysis was performed using Prism 5.0 (GraphPad Software). Statistical differences between two groups of data were determined using the unpaired two-tailed Student t-test. Data were considered significant when $p < 0.05$.

Results

KDM4D expression is elevated in GIST tissues

To dissect the contribution of demethylases to GIST, we first analysed the data from Oncomine. Surprisingly, we found that KDM4D levels were upregulated in GIST compared with normal tissues. To validate this finding, we further assessed KDM4D expression in our clinical GIST specimens. As shown in Fig. 1b, c and d, elevated KDM4D mRNA and protein expression was observed in GIST samples compared with matched normal tissues. Next, KDM4D expression was detected using IHC. Similarly, IHC results were consistent with prior observations



that KDM4D is preferentially overexpressed in GIST samples (Fig. 1e, f). These data indicate that KDM4D may be involved in GIST progression.

KDM4D promotes GIST cell proliferation, migration and invasion


Given the abnormal expression of KDM4D in GIST, we then assessed whether KDM4D expression affects GIST progression. To assess this hypothesis, we constructed the pcDNA-3.1(+)-KDM4D plasmid and control vector pcDNA-3.1(+) plasmid. Next, we transfected each plasmid into GIST-882 and GIST-T1 cells to obtain stable cell lines expressing KDM4D or control vector upon treatment with G418. As shown in Fig. 2a, western blot results revealed up-regulated KDM4D expression in GIST-882 and GIST-T1 cells expressing pcDNA-3.1(+)-KDM4D compared with vector cells. Further evaluating the role of KDM4D in GIST proliferation, we found, using CCK8 assays, that KDM4D cells exhibit increased growth compared with vector cells (Fig. 2b). Consistent with this result, BrdU/PI assay showed that overexpression of KDM4D increased the number of cells that incorporated BrdU compared with vector cells

(Fig. 2b), indicating that KDM4D cells exhibit increased DNA synthesis. Concordantly, KDM4D overexpression also increased the number and size of cell colonies, as determined by soft agar colony formation assays (Fig. 2d). These data indicate that KDM4D overexpression promotes GIST cell proliferation.

Clinically, approximately 20% of GISTs have metastasized at the time of diagnosis. Metastasis seriously limits the treatments available for GISTs. Thus, we further assessed whether KDM4D plays a role in GIST metastasis. As shown in Fig. 2e, overexpression KDM4D markedly increased cell migration in Transwell assays and promoted cell invasion through Matrigel. Consistent with this phenomenon, we observed that KDM4D overexpression strongly increased wound-healing capacity (Fig. 2f). Together, the above data suggest a role of KDM4D in GIST cell motility and invasion.

Silencing of KDM4D reduces GIST cell proliferation, migration and invasion

To better understand the role of KDM4D in GIST proliferation, migration and invasion, we performed




lentivirus-mediated knockdown using two independent shRNAs to silence KDM4D expression. Western blot revealed that the shRNAs effectively knocked down KDM4D in sh-KDM4D cells compared with scrambled shRNA (Fig. 3a). As shown in Fig. 3b, KDM4D knockdown strongly decreased GIST cell growth. In addition, KDM4D depletion also reduced DNA synthesis in both GIST-882 and GIST-T1 cells (Fig. 3c). We further found that silencing KDM4D markedly decreased the number and size of cell colonies (Fig. 3d) as determined by soft agar colony formation assays. Next, we assessed the role of KDM4D knockdown in GIST metastasis. Clearly, we

observed that KDM4D knockdown markedly abolished cell migration and invasion (Fig. 3e). Consistent with this result, KDM4D knockdown decreased cell mobility in a wound-healing assay (Fig. 3f). Taken together, these data suggest that silencing KDM4D actually inhibits GIST cell proliferation, migration and invasion.

The emerging role of KDM4D in GIST angiogenesis


Angiogenesis, which involves the formation of new capillaries from preexisting microvasculature, plays a vital role in tumour progression. Accumulating data indicate that tumour cells secrete several angiogenic factors (VEGF,



PDGFB, and bFGF) to promote angiogenesis [22]. The HIF/VEGFA signalling pathway plays a central role in tumour angiogenesis [23]. Thus, we further evaluate the function of KDM4D in tumour angiogenesis. Interestingly, KDM4D overexpression markedly up-regulated HIF1 β expression with minimal effects on HIF1A (which is mainly regulated by hypoxia) (Fig. 4a). In contrast, KDM4D knockdown attenuated HIF1 β expression (Fig. 4c). More importantly, elevated VEGFA expression was observed in KDM4D cells compared with control vector cells (Fig. 4a). Consistent with this result, downregulated VEGFA expression was detected in shKDM4D cells compared with shNC cells (Fig. 4c). In addition, KDM4D overexpression increased VEGFA secretion as confirmed using ELISA

assays (Fig. 4b). In contrast, silencing KDM4D reduced VEGFA secretion as confirmed using ELISA (Fig. 4d).

To further validate the role of KDM4D in angiogenesis, we cultured endothelial cells incubated with conditioned media (CM) from vector cells, KDM4D cells, shNC cells and shKDM4D cells, separately. As shown in Figs. 4 e and f, we found that CM from overexpression of KDM4D cells robustly induced tube formation of endothelial cells compared with CM from vector cells, and CM from knock-down KDM4D cells significantly decreased tube formation of endothelial cells compared with CM from shNC cells. Collectively, these data strongly suggest that KDM4D potentially influences angiogenesis through the Hif1 β /VEGFA signalling pathway.




HIF1 β transcriptional activation by KDM4D correlates closely with demethylation of H3K9me3 and H3K36me3

To further demonstrate whether VEGFA and HIF1 β overexpression is a mechanism regulated by KDM4D, we overexpressed KDM4D in shKDM4D cells. Western blot results showed that KDM4D rescue increased VEGFA and HIF1 β protein levels (Fig. 5a), indicating that the changes in VEGFA and HIF1 β are dependent on KDM4D. Then, we assessed whether KDM4D regulation of GIST phenotypes is dependent on the HIF1 β /VEGFA pathway. To assess this hypothesis, we transfected HIF1 β plasmids into ShKDM4D cells. CCK8 assays demonstrated that HIF1 β overexpression could

partially reverse the abolished proliferation mediated by knockdown of KDM4D (Additional file 1: Figure S1A). Furthermore, we found that HIF1 β overexpression reverses the attenuated tumour migration, invasion and angiogenesis mediated by silenced KDM4D (Additional file 1: Figure S1B, C). These data strongly indicate that the role of KDM4D in GIST proliferation, migration, invasion and angiogenesis is dependent on HIF1 β .

KDM4D is a member of the demethylase family and functions as a demethylase by mainly regulating the methylation levels of H3K36me3. Thus, to explore the mechanism of KDM4D-mediated regulation of HIF1 β expression, we first examined methylation levels of histone3 protein.



As shown in Figs. 5b, c, d and e, KDM4D overexpression decreased the methylation levels of both H3K36me3 and H3K9me3 compared with vector cells. In contrast, depletion of KDM4D strongly promoted H3K36me3 and H3K9me3 methylation, indicating that KDM4D might bind to the HIF1 β gene promoter to regulate its expression. To test this hypothesis, we examined the interaction between KDM4D protein and the HIF1 β gene promoter using CHIP assays. As shown in Fig. 5f, the HIF1 β gene promoter was directly immunoprecipitated with anti-KDM4D antibodies. Furthermore, CHIP assay

results revealed that KDM4D overexpression decreased the interaction between the HIF1 β gene promoter and both H3K36me3 and H3K9me3 compared with vector cells and that silencing KDM4D enhanced the interaction (Fig. 5g, i). Next, we further determined whether KDM4D activates the HIF1 β gene promoter. Using dual-luciferase assays, we found, interestingly, that HIF1 β gene promoter activity was activated in KDM4D cells compared with control vector cells (Fig. 5h). In contrast, KDM4D significantly attenuated Hif1 β gene promoter activity compared with shNC cells (Fig. 5j). Taken together, our results

demonstrate that KDM4D transcriptionally activates HIF1 β expression via demethylation of H3K9me3 and H3K36me3 at the promoter region.


KDM4D overexpression promotes GIST cell proliferation and angiogenesis in vivo

To examine the in vivo pro-tumour activity by KDM4D, vector or stably expressed KDM4D cells were injected into right subcutaneous tissues of Balb/c nude mice. As shown in Fig. 6a, vector cell growth was significantly retarded compared with KDM4D cells in mice. In addition, the weight and size of tumours formed from vector cells were decreased compared with tumours formed from KDM4D cells (Fig. 6b, c). Importantly, immunohistochemistry (IHC) assay results also demonstrated that CD31 expression was significantly upregulated in the KDM4D group compared with the control vector group, indicating that tumour microvessel density was strongly increased in the KDM4D group (Fig. 6d, e).

Consistent with this finding, western blot results also demonstrated VEGFA expression was strongly increased in the KDM4D group compared with the control vector group (Fig. 6f).

KDM4D knockdown suppresses GIST cell proliferation and angiogenesis in vivo

To further demonstrate the function of KDM4D in GIST proliferation and angiogenesis, ShNC or stably expressing ShKDM4D cells were injected into the right subcutaneous tissues of Balb/c nude mice, separately. In contrast to the overexpression tumour xenograft model, we found, interestingly, that ShNC cells grew significantly faster compared with ShKDM4D cells in mice (Fig. 7a). In addition, the weight and size of tumours formed from ShNC cells were increased compared with tumours formed from ShKDM4D cells (Fig. 7b, c). To assess the status of angiogenesis in xenograft tumours, we examined CD31 expression in the three groups using IHC assays. As shown in






Fig. 7d and e, CD31 expression was downregulated in the ShKDM4D group compared with the ShNC group, indicating that the microvessel density was attenuated because KDM4D was silenced in GISTs. Next, we examined the relationship among KDM4D, VEGFA and HIF1 β in xenograft mouse models. Western blot results clearly demonstrated that VEGFA and HIF1 β expression was strongly decreased in the ShKDM4D group compared with the ShNC group (Fig. 7f). Collectively, these data indicate that KDM4D plays a crucial role in GIST proliferation and angiogenesis.

Discussion

Most GISTs harbouring mutant KIT or PDGFRA are treated with targeted therapies, such as imatinib. Although limited success is noted at the beginning of treatment, the majority of patients ultimately develop drug resistance [24]. More importantly, wild type GISTs do not respond to molecular targeted therapy [25–29].

Thus, further understanding of the molecular mechanism regulating GIST progression is urgent. Novel treatment strategies should be identified to treat patients with GIST, especially those with unresectable and advanced GIST.

In this study, we found that KDM4D is overexpressed in GISTs compared with corresponding normal tissues. This finding suggests that epigenetic post-translational modifications may be involved in GIST progression. Consistent with our findings, previous data also demonstrated that epigenetic alterations play important roles in GIST development and even have predictive value in GISTs. For example, CD133 methylation percentage is inversely correlated with CD133 protein expression, and hypermethylation in small (< 2 cm) GIST reflects tumour size. Niimura et al. demonstrated that multiple genes in the HOXC cluster are marked by active histone H3K4me3 in malignant GISTs and revealed that HOTAIR might be a potentially useful biomarker in malignant GIST [30]. These data suggested that histone-modifying enzymes could act as novel

therapeutic targets in GISTs. KDM4D expression in GISTs has not been previously assessed. Herein, we first evaluated the roles of KDM4D in GIST cell proliferation, migration, and invasion. Surprisingly, KDM4D overexpression strongly increased GIST cell proliferation, migration, and invasion. In contrast, KDM4D depletion significantly attenuated GIST cell proliferation, migration, and invasion, indicating that KDM4D regulated GIST progression. Targeting KDM4D might offer an effective method to control the development of GISTs.

Angiogenesis is a complicated process that is crucial for cancer proliferation and metastasis [31]. Evidence indicates that the Hif/VEGFA signalling pathway is a potent driver of angiogenesis [32]. In the present study, we evaluated the role of KDM4D in tumour angiogenesis. We found that KDM4D overexpression strongly accelerates HUVEC tube formation both in vitro and in vivo. In addition, increased VEGFA secretion was observed in KDM4D cells compared with control vector cells. Consistent with the overexpression experiments, we further found that silencing KDM4D significantly abolished tube formation and decreased VEGFA secretion, indicating that KDM4D regulates GIST development in an angiogenesis-dependent manner. The Hif family plays a critical role in angiogenesis, especially HIF1 α , which is mainly regulated by hypoxia. Other Hif members also cooperate with HIF1 α to play an important role in tumour progression, including tumour angiogenesis. HIF1 β also regulates angiogenesis in a HIF1 α -independent manner [33]. To explain the underlying molecular mechanisms of KDM4D-induced angiogenesis, we examined HIF1 α and HIF1 β protein expression. Interestingly, KDM4D markedly upregulated HIF1 β expression but not HIF1 α expression.

KDM4D regulates gene expression by regulating the methylation levels of both H3K36me3 and H3K9me3 [34, 35]. Consistent with previous reports, KDM4D overexpression significantly increased H3K36me3 and H3K9me3 methylation levels compared with vector cells. Importantly, CHIP assays revealed that KDM4D interacts with the HIF1 β promoter. In addition, changes in KDM4D strongly affect the interaction between the HIF1 β promoter and both H3K36me3 and H3K9me3. These findings indicated that KDM4D induces tumour angiogenesis via HIF1 β expression. In addition, compared with the control vector group in Balb/c nude mice, the growth of KDM4D-overexpressing cells was increased along with increased VEGFA expression as assessed by immunohistochemistry. These results indicate the pro-tumour effect of KDM4D via tumour angiogenesis.

In summary, our study presents the following lines of evidence supporting the notion that KDM4D plays a vital role in GIST progression. First, KDM4D expression is upregulated in GIST samples. Second, we demonstrated that KDM4D plays a central role in GIST proliferation,

migration and invasion both in vitro and in vivo. Third, we demonstrated that KDM4D is required for tumour angiogenesis via regulating VEGFA secretion. Finally, we further revealed that KDM4D directly binds to and activates the HIF1 β gene promoter.

Conclusions

These results identify a novel regulatory pathway (KDM4D/HIF1 β /VEGFA) involved in GIST progression. Thus, we propose that elevated KDM4D expression may serve as a biomarker and act as a novel potential target for GIST.

Additional file

Additional file 1: Figure S1A. GIST882 cells stably expressing ShNC or ShKDM4D were transfected with HIF1 β plasmids or control plasmids, after transfection for 24h, then, the indicated GIST882 cells were plated into 96-wells. Then, the cell viability of the indicated GIST882 cells was detected at the indicated time by CCK8 assay. **Figure S1B.** Cells were plated in the upper chamber with/without Matrigel for 24 h. Then, cell migrative and invasive ability was assessed by observing the cells migrating to the underside of transwell insert. **Figure S1C.** After transfection for 48h, conditional medium from the indicated GIST882 cells were used to incubated with HUVECs cells in 96 well plates. The tube formation was examined. (TIF 2962 kb)

Abbreviations

GIST: Gastrointestinal stromal tumour; HIF1 β : Aryl hydrocarbon receptor nuclear translocator; IHC staining: Immunohistochemical staining; KDM4D: Lysine-specific demethylase 4D; ShRNA: Short hairpin RNA; VEGFA: Vascular endothelial growth factor A

Funding

This work was supported by grants from the Science and Technology of Hubei Province of China (No. 2016CFB136) and the Science and Technology of Hubei Province of China (No. 2016CFB550).

Availability of data and materials

All data generated or analysed during this study are included in this published article.

Authors' contributions

XL, JH and XY designed this research. FH performed most of the experiments, analysed the data, generated the figures and drafted this manuscript. HL and LL performed western blot experiments, qRT-PCR and CCK-8 assays. FX and SL helped with luciferase, colony formation, tube formation, CCK-8, BrdU/PI and Transwell assays. All authors read and approved the final manuscript.

Ethics approval and consent to participate

GIST tissues were obtained from patients who underwent surgical resection at Tongji Hospital (Wuhan, China). Consent letters were obtained from patients before this study. All experiments were approved by the Ethics Committee of Tongji Hospital and conducted in accordance with the Code of Ethics of the World Medical Association (Declaration of Helsinki).

Consent for publication

We obtained consent to publish this paper from all study participants.

Competing interests

The authors declare that they have no competing interests.

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Received: 20 April 2018 Accepted: 23 July 2018

Published online: 30 July 2018

References

- Corless CL, Barnett CM, Heinrich MC. Gastrointestinal stromal tumours: origin and molecular oncology. *Nat Rev Cancer*. 2011;11(12):865–78.
- Niinuma T, Suzuki H, Sugai T. Molecular characterization and pathogenesis of gastrointestinal stromal tumor. *Transl Gastroenterol Hepatol*. 2018;3:2.
- Schaefer IM, Marino-Enriquez A, Fletcher JA. What is new in gastrointestinal stromal tumor? *Adv Anat Pathol*. 2017;24(5):259–67.
- Joensuu H, Hohenberger P, Corless CL. Gastrointestinal stromal tumour. *Lancet*. 2013;382(9896):973–83.
- Joensuu H, Wardelmann E, Sihto H, Eriksson M, Sundby Hall K, Reichardt A, Hartmann JT, Pink D, Cameron S, Hohenberger P, Al-Batran SE, Schlemmer M, Bauer S, Nilsson B, Kallio R, Junnila J, Vehtari A, Reichardt P. Effect of KIT and PDGFRA mutations on survival in patients with gastrointestinal stromal tumors treated with adjuvant Imatinib: an exploratory analysis of a randomized clinical trial. *JAMA Oncol*. 2017;3(5):602–9.
- Tamborini E. Mechanism of resistance in gastrointestinal stromal tumors. *Handb Exp Pharmacol*. 2017; https://doi.org/10.1007/164_2017_74.
- Kang YK, Kang BW, Im SA, Lee JL, Park SR, Kang WK, Chang HM, Kim TW, Oh DY, Jung KH, Ryu MH. Two-year adjuvant imatinib mesylate after complete resection of localized, high-risk GIST with KIT exon 11 mutation. *Cancer Chemother Pharmacol*. 2013;71(1):43–51.
- Huss S, Elges S, Trautmann M, Sperveslage J, Hartmann W, Wardelmann E. Classification of KIT/PDGFRA wild-type gastrointestinal stromal tumors: implications for therapy. *Expert Rev Anticancer Ther*. 2015;15(6):623–8.
- Jones PA, Baylin SB. The epigenomics of cancer. *Cell*. 2007;128(4):683–92.
- Wainwright EN, Scaffidi P. Epigenetics and Cancer stem cells: unleashing, hijacking, and restricting cellular plasticity. *Trends Cancer*. 2017;3(5):372–86.
- Gedder H, Braun A, Kayser C, Dimmler A, Faller G, Agaimy A, Haller F, Moskalev EA. Epigenetic regulation of CD133 in gastrointestinal stromal tumors. *Am J Clin Pathol*. 2017;147(5):515–24.
- Igarashi S, Suzuki H, Niinuma T, Shimizu H, Nojima M, Iwaki H, Nobuoka T, Nishida T, Miyazaki Y, Takamaru H, Yamamoto E, Yamamoto H, Tokino T, Hasegawa T, Hirata K, Imai K, Toyota M, Shinomura Y. A novel correlation between LINE-1 hypomethylation and the malignancy of gastrointestinal stromal tumors. *Clin Cancer Res*. 2010;16(21):5114–23.
- Abdelfatah E, Kerner Z, Nanda N, Ahuja N. Epigenetic therapy in gastrointestinal cancer: the right combination. *Therap Adv Gastroenterol*. 2016;9(4):560–79.
- Huang KK, McPherson JR, Tay ST, Das K, Tan IB, Ng CC, Chia NY, Zhang SL, Myint SS, Hu L, Rajasegaran V, Huang D, Loh JL, Gan A, Sairi AN, Sam XX, Dominguez LT, Lee M, Soo KC, Ooi LL, Ong HS, Chung A, Chow PK, Wong WK, Selvarajan S, Ong CK, Lim KH, Nandi T, Rozen S, Teh BT, Quek R, Tan P. SETD2 histone modifier loss in aggressive GI stromal tumours. *Gut*. 2016; 65(12):1960–72.
- Huang T, Lin C, Zhong LL, Zhao L, Zhang G, Lu A, Wu J, Bian Z. Targeting histone methylation for colorectal cancer. *Therap Adv Gastroenterol*. 2017;10(1):114–31.
- Liu Y, Tseng M, Perdreau SA, Rossi F, Antonescu C, Besmer P, Fletcher JA, Duensing S, Duensing A. Histone H2AX is a mediator of gastrointestinal stromal tumor cell apoptosis following treatment with imatinib mesylate. *Cancer Res*. 2007;67(6):2685–92.
- Metzger E, Stepputtis SS, Strietz J, Preca BT, Urban S, Willmann D, Allen A, Zenk F, Iovino N, Bronsert P, Proske A, Follo M, Boerries M, Stickeler E, Xu J, Wallace MB, Stafford JA, Kanouni T, Maurer J, Schule R. KDM4 inhibition targets breast Cancer stem-like cells. *Cancer Res*. 2017;77(21):5900–12.
- Agger K, Miyagi S, Pedersen MT, Kooistra SM, Johansen JV, Helin K. Jmjd2/Kdm4 demethylases are required for expression of Il3ra and survival of acute myeloid leukemia cells. *Genes Dev*. 2016;30(11):1278–88.
- Awwad SW, Ayoub N. Overexpression of KDM4 lysine demethylases disrupts the integrity of the DNA mismatch repair pathway. *Biol Open*. 2015;4(4):498–504.
- Li H, Yang X, Wang G, Li X, Tao D, Hu J, Luo X. KDM4B plays an important role in mitochondrial apoptosis by upregulating HAX1 expression in colorectal cancer. *Oncotarget*. 2016;7(36):57866–77.
- Wang G, Cao X, Lai S, Luo X, Feng Y, Xia X, Yen PM, Gong J, Hu J. PI3K stimulates DNA synthesis and cell-cycle progression via its p55PIK regulatory subunit interaction with PCNA. *Mol Cancer Ther*. 2013;12(10):2100–9.
- Guo P, Wang Y, Dai C, Tao C, Wu F, Xie X, Yu H, Zhu Q, Li J, Ye L, Yu F, Shan Y, Yu Z, Dhanasekaran R, Zheng R, Chen G. Ribosomal protein S15a promotes tumor angiogenesis via enhancing Wnt/beta-catenin-induced FGF18 expression in hepatocellular carcinoma. *Oncogene*. 2018;37(9):1220–36.
- Weijts BG, Bakker WJ, Cornelissen PW, Liang KH, Schaftenaar FH, Westendorp B, de Wolf CA, Paciejewska M, Scheele CL, Kent L, Leone G, Schulte-Merker S, de Bruin A. E2F7 and E2F8 promote angiogenesis through transcriptional activation of VEGFA in cooperation with HIF1. *EMBO J*. 2012;31(19):3871–84.
- Joensuu H, Eriksson M, Sundby Hall K, Reichardt A, Hartmann JT, Pink D, Ramadori G, Hohenberger P, Al-Batran SE, Schlemmer M, Bauer S, Wardelmann E, Nilsson B, Sihto H, Bono P, Kallio R, Junnila J, Alvegard T, Reichardt P. Adjuvant Imatinib for high-risk GI stromal tumor: analysis of a randomized trial. *J Clin Oncol*. 2016;34(3):244–50.
- Bauer S, Parry JA, Muhlenberg T, Brown MF, Seneviratne D, Chatterjee P, Chin A, Rubin BP, Kuan SF, Fletcher JA, Duensing S, Duensing A. Proapoptotic activity of bortezomib in gastrointestinal stromal tumor cells. *Cancer Res*. 2010;70(1):150–9.
- Casali PG, Le Cesne A, Poveda Velasco A, Kotasek D, Rutkowski P, Hohenberger P, Fumagalli E, Judson IR, Italiano A, Gelderblom H, Adenis A, Hartmann JT, Duffaud F, Goldstein D, Broto JM, Gronchi A, Dei Tos AP, Marraud S, van der Graaf WT, Zalcberg JR, Litiere S, Blay JY. Time to Definitive Failure to the First Tyrosine Kinase Inhibitor in Localized GI Stromal Tumors Treated With Imatinib As an Adjuvant: A European Organisation for Research and Treatment of Cancer Soft Tissue and Bone Sarcoma Group Intergroup Randomized Trial in Collaboration With the Australasian Gastro-Intestinal Trials Group, UNICANCER, French Sarcoma Group, Italian Sarcoma Group, and Spanish Group for Research on Sarcomas. *J Clin Oncol*. 2015;33(36):4276–83.
- Emile JF, Brahimi S, Coindre JM, Bringuiet PP, Monges G, Samb P, Doucet L, Hostein I, Landi B, Buisine MP, Neuville A, Bouche O, Cervera P, Pretet JL, Tisserand J, Gauthier A, Le Cesne A, Sabourin JC, Scaozec JY, Bonvalot S, Corless CL, Heinrich MC, Blay JY, Aegerter P. Frequencies of KIT and PDGFRA mutations in the MolecGIST prospective population-based study differ from those of advanced GISTs. *Med Oncol*. 2012;29(3):1765–72.
- Joensuu H, Eriksson M, Sundby Hall K, Hartmann JT, Pink D, Schutte J, Ramadori G, Hohenberger P, Duyster J, Al-Batran SE, Schlemmer M, Bauer S, Wardelmann E, Sarlomo-Rikala M, Nilsson B, Sihto H, Monge OR, Bono P, Kallio R, Vehtari A, Leinonen M, Alvegard T, Reichardt P. One vs three years of adjuvant imatinib for operable gastrointestinal stromal tumor: a randomized trial. *Jama*. 2012;307(12):1265–72.
- Medeiros F, Corless CL, Duensing A, Hornick JL, Oliveira AM, Heinrich MC, Fletcher JA, Fletcher CD. KIT-negative gastrointestinal stromal tumors: proof of concept and therapeutic implications. *Am J Surg Pathol*. 2004;28(7):889–94.
- Niinuma T, Suzuki H, Nojima M, Noshio K, Yamamoto H, Takamaru H, Yamamoto E, Maruyama R, Nobuoka T, Miyazaki Y, Nishida T, Bamba T, Kanda T, Ajioka Y, Taguchi T, Okahara S, Takahashi H, Nishida Y, Hosokawa M, Hasegawa T, Tokino T, Hirata K, Imai K, Toyota M, Shinomura Y. Upregulation of miR-196a and HOTAIR drive malignant character in gastrointestinal stromal tumors. *Cancer Res*. 2012;72(5):1126–36.
- Carmeliet P, Jain RK. Molecular mechanisms and clinical applications of angiogenesis. *Nature*. 2011;473(7347):298–307.
- Potente M, Gerhardt H, Carmeliet P. Basic and therapeutic aspects of angiogenesis. *Cell*. 2011;146(6):873–87.
- Wondimu A, Weir L, Robertson D, Mezentsev A, Kalachikov S, Panteleyev AA. Loss of Arnt (Hif1beta) in mouse epidermis triggers dermal angiogenesis, blood vessel dilation and clotting defects. *Lab Invest*. 2012;92(1):110–24.
- Wu R, Wang Z, Zhang H, Gan H, Zhang Z. H3K9me3 demethylase Kdm4d facilitates the formation of pre-initiative complex and regulates DNA replication. *Nucleic Acids Res*. 2017;45(1):169–80.
- Zoabi M, Nadar-Ponniath PT, Khoury-Haddad H, Usaj M, Budowski-Tal I, Haran T, Henn A, Mandel-Gutfreund Y, Ayoub N. RNA-dependent chromatin localization of KDM4D lysine demethylase promotes H3K9me3 demethylation. *Nucleic Acids Res*. 2014;42(21):3026–38.